



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

the symbols in (5) occur on this triangle in the direction contrary to that of the hands of a watch, and the displacement of the letters takes place in this direction; thus

$$CRx = \begin{matrix} A & B \\ P & Q \end{matrix}$$

On the other hand

$$RCx = \begin{matrix} B & P \\ A & Q \end{matrix}$$

11. In comparing any two of the 24 square arrangements of the four letters, one of the letters may by the process indicated in equation (2) be brought into the same position in each form; the forms will then differ either in the position of two of the remaining letters indicating one of the relations denoted by R , C or J or else in the position of all three letters indicating one of the relations denoted by CR or RC , discussed above.

SOLUTION OF PROB. 407 BY PROF. EDGAR FRISBY.—I notice that the solution of problem 407 [see p. 158] contains a remarkable mistake in omitting the exponent $\frac{1}{2}$. The expression is an elliptic integral; it can be reduced to a series thus:

$$\int_0^1 \frac{(1+x^4)^{\frac{1}{2}}}{(1-x)^{\frac{1}{2}}} dx = \int_0^1 \frac{1 + \frac{1}{2}x^4 - \frac{1}{2} \cdot \frac{1}{4}x^8 + \frac{1}{2} \cdot \frac{1}{4} \cdot \frac{3}{6}x^{12} - \frac{1}{2} \cdot \frac{1}{4} \cdot \frac{3}{6} \cdot \frac{5}{8}x^{16} + \dots}{\sqrt{1-x^2}} dx.$$

Integrating each term by parts and only retaining the last term of each series, as the other terms vanish at both limits, we have

$$\int_0^{\frac{\pi}{2}} (1+\cos^4\theta)^{\frac{1}{2}} d\theta = \frac{\pi}{2} \left\{ \frac{1 \cdot 1 \cdot 3}{2 \cdot 2 \cdot 4} - \frac{1 \cdot 1 \cdot 1 \cdot 3 \cdot 5 \cdot 7}{2 \cdot 4 \cdot 2 \cdot 4 \cdot 6 \cdot 8} + \frac{1 \cdot 1 \cdot 1 \cdot 3 \cdot 5 \cdot 7 \cdot 9 \cdot 11}{2 \cdot 4 \cdot 6 \cdot 2 \cdot 4 \cdot 6 \cdot 8 \cdot 10 \cdot 12 \dots} \right\}.$$

SOLUTIONS OF PROBLEMS IN NUMBER FIVE.

SOLUTIONS of problems in No. 5 have been received as follows:

From Prof. L. G. Barbour, 412; Alex. S. Christie, 411; Prof. W. P. Casey, 409, 412, 416; Geo. Eastwood, 413, 416, 417; C. E. Everett, 409, 410, 415, 416; Prof. Edgar Frisby, 409, 411, 415; Wm. Hoover, 409, 411, 413, 416; L. S. Hulbert, 412; E. H. Moore, Jr., 416; Prof. P. H. Philbrick, 409, 414, 415, 416; A. L. Parman, 409; Prof. E. B. Seitz, 409; Prof. J. Scheffer, 409, 410, 411, 416; Prof. C. M. Woodward, 409, 410, 413; R. S. Woodward, 413.

[Solutions of 401, 402, 403 and 406 by Chas. E. Everett and Prof. Casey, respectively, were received too late for acknowledgment in No. 5.]